



Pain Control: Adult



History


- Age
- Location
- Duration
- Severity (1 - 10)
- If child use Wong-Baker faces scale
- Past medical history
- Medications
- Drug allergies

Signs and Symptoms

- Severity (pain scale)
- Quality (sharp, dull, etc.)
- Radiation
- Relation to movement, respiration
- Increased with palpation of area

Differential

- Per the specific protocol
- Musculoskeletal
- Visceral (abdominal)
- Cardiac
- Pleural / Respiratory
- Neurogenic
- Renal (colic)



General Protocols

Legend		
	MR	
B	EMT	B
I	EMT- I	I
P	EMT- P	P
M	Medical Control	M

Pearls

- **Recommended Exam: Mental Status, Area of Pain, Neuro**
- Pain severity (0-10) is a vital sign to be recorded pre and post IV or IM medication delivery and at disposition.
- Vital signs should be obtained pre, 15 minutes post, and at disposition with all pain medications.
- Patients with presumed kidney stone should first receive Toradol. A narcotic may then be considered.
- Contraindications to the use of a **narcotic** include hypotension, head injury, respiratory distress or severe COPD.
- **Ketorolac (Toradol) and Ibuprofen should not be used in patients with known renal disease or renal transplant, in patients who have known drug allergies to NSAID's (non-steroidal anti-inflammatory medications), with active bleeding, or in patients who may need surgical intervention such as open fractures or fracture deformities.**
- All patients should have drug allergies documented prior to administering pain medications.
- All patients who receive IM or IV medications must be observed 15 minutes for drug reaction.
- **Ibuprofen or Ketorolac** should not be given for headaches or abdominal pain, history of gastritis, stomach ulcers, fracture, or if patient will require sedation
- Do not administer any PO medications for patients who may need surgical intervention such as open fractures or fracture deformities, headaches, or abdominal pain.
- Do not administer **Acetaminophen** to patients with a history of liver disease.
- See drug list for other contraindications for Narcotics, Acetaminophen, Nitrous Oxide, Ketorolac, and Ibuprofen.