

Police Custody



History


- Traumatic Injury
- Drug Abuse
- Cardiac History
- History of Asthma
- Psychiatric History

Signs and Symptoms

- External signs of trauma
- Palpitations
- Shortness of breath
- Wheezing
- Altered Mental Status
- Intoxication/Substance Abuse

Differential

- Agitated Delirium Secondary to Psychiatric Illness
- Agitated Delirium Secondary to Substance Abuse
- Traumatic Injury
- Closed Head Injury
- Asthma Exacerbation
- Cardiac Dysrhythmia



Pearls

- For this protocol to be used, the patient does not have to be under police custody.
- Agitated delirium is characterized by marked restlessness, irritability, and/or high fever. Patients exhibiting these signs are at high risk for sudden death and should be transported to hospital by ALS personnel.
- Patients restrained by law enforcement devices cannot be transported in the ambulance without a law enforcement officer in the patient compartment who is capable of removing the devices.
- If there is any doubt about the cause of a patient's alteration in mental status, transport the patient to the hospital for evaluation.
- If an asthmatic patient is exposed to pepper spray and released to law enforcement, all parties should be advised to immediately recontact EMS if wheezing/difficulty breathing occurs.
- All patients in police custody retain the right to request transport. This should be coordinated with law enforcement.
- If extremity/chemical/law enforcement restraints are applied, completed Restraint procedure in call reporting system.

Protocol 14

Any local EMS System changes to this document must follow the NC OEMS Protocol Change Policy and be approved by OEMS

2009